



Mylan
Quality
Pharmaceutical
Products



Mylan Pharmaceuticals Inc.
781 Chestnut Ridge Road
Morgantown, WV 26504-4310
Telephone: 800-796-9526
Fax: 304-598-3232
www.mylan.com

Mylan

MYCOPHENOLATE MOFETIL CAPSULES

New Product Announcement
MYLAN PHARMACEUTICALS INTRODUCES
Mycophenolate Mofetil Capsules, 250 mg[†]

Compare To: CellCept[®] Capsules
Therapeutic Rating: AB
Expiration Dating: 24 Months
ANDA Number: 65-520
Product Category: Immunosuppressives

[†] Please See Attached Full Prescribing Information including, Black Box Warning and Medication Guide.

Strength:	250 mg	
Form:	Capsules	
Color (Cap):	Caramel Opaque	
Color (Body):	Lavender Opaque	
Imprint (Cap):	MYLAN over 2250	
Imprint (Body):	MYLAN over 2250	
Package Size:	100s	500s
NDC #:	0378-2250-01	0378-2250-05
Bottle Size:	120 cc	18 oz.
Case Pack:	144	12
Inner Pack:	12 x 12 x 100	12 x 1 x 500
Length:	16 1/2	17 5/8
Width:	11 3/8	12 1/2
Height:	12 3/8	5 3/8
Cube:	1.48	0.49
Weight:	18.72 lbs.	6.36 lbs.
AWP[‡]:	\$396.24	\$1,981.20

[‡]AWP for a Mylan product is reported by Mylan with reference to AWP for a brand company's therapeutically-equivalent product, as reported by American Druggist, First Data Bank or another nationally recognized publication. AWP reported by Mylan, however, is not necessarily the same as the AWP that might be independently established and reported by the publisher. AWP does not take into account any discounts, chargebacks, rebates, or other reductions in price that may be provided. AWP should not be relied upon as the actual cost to the pharmacy or to the customer or consumer. AWP is subject to change at any time.

MYCOPHENOLATE MOFETIL CAPSULES, 250 mg and MYCOPHENOLATE MOFETIL TABLETS, 500 mg


Rx only

WARNING: Immunosuppression may lead to increased susceptibility to infection and possible development of lymphoma. Only physicians experienced in immunosuppressive therapy and management of renal, cardiac or hepatic transplant patients should use mycophenolate mofetil. Patients receiving the drug should be managed in facilities equipped and staffed with adequate laboratory and supportive medical resources. The physician responsible for maintenance therapy should have complete information requisite for the follow-up of the patient.

Female users of childbearing potential must use contraception. Use of mycophenolate mofetil during pregnancy is associated with increased risk of pregnancy loss and congenital malformations.

DESCRIPTION: Mycophenolate mofetil is the 2-morpholinoethyl ester of mycophenolic acid (MPA), an immunosuppressive agent; inosine monophosphate dehydrogenase (IMPDH) inhibitor.

The chemical name for mycophenolate mofetil is 2-Morpholinoethyl (E)-6-(4-hydroxy-6-methoxy-7-methyl-3-oxo-5-phthalanyl)-4-methyl-4-hexenoate. It has a molecular formula of $C_{23}H_{31}NO_7$, a molecular weight of 433.5, and the following structural formula:



Mycophenolate mofetil is a white to almost white crystalline powder. It is practically insoluble in water (43 mcg/mL at pH 7.4), the solubility increases in acidic medium (4.27 mg/mL at pH 3.6). It is freely soluble in acetone, soluble in methanol, and sparingly soluble in anhydrous ethanol. The apparent partition coefficient in 1-octanol/water (pH 7.4) buffer solution is 238. The pKa values for mycophenolate mofetil are 5.6 for the morpholino group and 8.5 for the phenolic group.

Mycophenolate mofetil is available for oral administration as capsules containing 250 mg of mycophenolate mofetil and tablets containing 500 mg of mycophenolate mofetil.

Mycophenolate Mofetil Capsules, 250 mg contain the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, pregelatinized starch, and sodium lauryl sulfate. The empty gelatin capsule shells contain black iron oxide, FD&C Blue No. 2, gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. In addition, the imprinting ink contains the following: ammonium hydroxide, black iron oxide, propylene glycol, and shellac glaze.

Mycophenolate Mofetil Tablets, 500 mg contain the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, red iron oxide, sodium lauryl sulfate, talc, titanium dioxide and yellow iron oxide.

CLINICAL PHARMACOLOGY: Mechanism of Action: Mycophenolate mofetil has been demonstrated in experimental animal models to prolong the survival of allogeneic transplants (kidney, heart, liver, intestine, limb, small bowel, pancreatic islets and bone marrow).

Mycophenolate mofetil has also been shown to reverse ongoing acute rejection in the canine renal and rat cardiac allograft models. Mycophenolate mofetil also inhibited proliferative arteriopathy in experimental models of aortic and cardiac allografts in rats, as well as in primate cardiac xenografts. Mycophenolate mofetil was used alone or in combination with other immunosuppressive agents in these studies. Mycophenolate mofetil has been demonstrated to inhibit immunologically mediated inflammatory responses in animal models and to inhibit tumor development and prolong survival in murine tumor transplant models.

Mycophenolate mofetil is rapidly absorbed following oral administration and hydrolyzed to form MPA, which is the active metabolite. MPA is a potent, selective, uncompetitive and reversible inhibitor of inosine monophosphate dehydrogenase (IMPDH), and therefore inhibits the *de novo* pathway of guanosine nucleotide synthesis without incorporation into DNA. Because T- and B-lymphocytes are critically dependent for their proliferation on *de novo* synthesis of purines, whereas other cell types can utilize salvage pathways, MPA has potent cytostatic effects on lymphocytes. MPA inhibits proliferative responses of T- and B-lymphocytes to both mitogenic and allospecific stimulation. Addition of guanosine or deoxyguanosine reverses the cytostatic effects of MPA on lymphocytes. MPA also suppresses antibody formation by B-lymphocytes. MPA prevents the glycosylation of lymphocyte and monocyte glycoproteins that are involved in intercellular adhesion to endothelial cells and may inhibit recruitment of leukocytes into sites of inflammation and graft rejection. Mycophenolate mofetil did not inhibit early events in the activation of human peripheral blood mononuclear cells, such as the production of interleukin-1 (IL-1) and interleukin-2 (IL-2), but did block the coupling of these events to DNA synthesis and proliferation.

Pharmacokinetics: Following oral and intravenous administration, mycophenolate mofetil undergoes rapid and complete metabolism to MPA, the active metabolite. Oral absorption of the drug is rapid and essentially complete. MPA is metabolized to form the phenolic glucuronide of MPA (MPAG) which is not pharmacologically active. The parent drug, mycophenolate mofetil, can be measured systemically during the intravenous infusion; however, shortly (about 5 minutes) after the infusion is stopped or after oral administration, mycophenolate mofetil concentration is below the limit of quantitation (0.4 mcg/mL).

Absorption: In 12 healthy volunteers, the mean absolute bioavailability of oral mycophenolate mofetil relative to intravenous mycophenolate mofetil (based on MPA AUC) was 94%. The area under the plasma-concentration time curve (AUC) for MPA appears to increase in a dose proportional fashion in renal transplant patients receiving multiple doses of mycophenolate mofetil up to a daily dose of 3 g (see Table 1).

Food (27 g fat, 650 calories) had no effect on the extent of absorption (MPA AUC) of mycophenolate mofetil when administered at doses of 1.5 g bid to renal transplant patients. However, MPA C_{max} was decreased by 40% in the presence of food (see DOSAGE AND ADMINISTRATION).

Distribution: The mean (\pm SD) apparent volume of distribution of MPA in 12 healthy volunteers is approximately 3.6 (\pm 1.5) and 4 (\pm 1.2) L/kg following intravenous and oral administration, respectively. MPA, at clinically relevant concentrations, is 97% bound to plasma albumin. MPAG is 82% bound to plasma albumin at MPAG concentration ranges that are normally seen in stable renal transplant patients; however, at higher MPAG concentrations (observed in patients with renal impairment or delayed renal graft function), the binding of MPA may be reduced as a result of competition between MPAG and MPA for protein binding. Mean blood to plasma ratio of radioactivity concentrations was approximately 0.6 indicating that MPA and MPAG do not extensively distribute into the cellular fractions of blood.

In vitro studies to evaluate the effect of other agents on the binding of MPA to human serum albumin (HSA) or plasma proteins showed that salicylate (at 25 mg/dL with HSA) and MPAG (at \approx 460 mcg/mL with plasma proteins) increased the free fraction of MPA. At concentrations that exceeded what is encountered clinically, cyclosporine, digoxin, naproxen, prednisone, propranolol, tacrolimus, theophylline, tolbutamide and warfarin did not increase the free fraction of MPA. MPA at concentrations as high as 100 mcg/mL had little effect on the binding of warfarin, digoxin or propranolol, but decreased the binding of theophylline from 53% to 45% and pentylenol from 90% to 87%.

Metabolism: Following oral and intravenous dosing, mycophenolate mofetil undergoes complete metabolism to MPA, the active metabolite. Metabolism to MPA occurs presystemically after oral dosing. MPA is metabolized principally by glucuronyl transferase to form the phenolic glucuronide of MPA (MPAG) which is not pharmacologically active. *In vivo*, MPAG is converted to MPA via enterohepatic recirculation. The following metabolites of the 2-hydroxyethyl-morpholino moiety are also recovered in the urine following oral administration of mycophenolate mofetil to healthy subjects: N-(2-carboxymethyl)-morpholine, N-(2-hydroxyethyl)-morpholine, and the N-oxide of N-(2-hydroxyethyl)-morpholine.

Secondary peaks in the plasma MPA concentration-time profile are usually observed 6 to 12 hours post-dose. The coadministration of cholestyramine (4 g tid) resulted in approximately a 40% decrease in the MPA AUC (largely as a consequence of lower concentrations in the terminal portion of the profile). These observations suggest that enterohepatic recirculation contributes to MPA plasma concentrations.

Increased plasma concentrations of mycophenolate mofetil metabolites (MPA 50% increase and MPAG about a 3-fold to 6-fold increase) are observed in patients with renal insufficiency (see CLINICAL PHARMACOLOGY: Special Populations).

Excretion: Negligible amount of drug is excreted as MPA (< 1% of dose) in the urine. Orally administered radiolabeled mycophenolate mofetil resulted in complete recovery of the administered dose, with 93% of the administered dose recovered in the urine and 6% recovered in feces. Most (about 87%) of the administered dose is excreted in the urine as MPAG. At clinically encountered concentrations, MPA and MPAG are usually not removed by hemodialysis. However, at high MPAG plasma concentrations (> 100 mcg/mL), small amounts of MPAG are removed. Bile acid sequestrants, such as cholestyramine, reduce MPA AUC by interfering with enterohepatic circulation of the drug (see OVERDOSAGE).

Mean (\pm SD) apparent half-life and plasma clearance of MPA are 17.9 (\pm 6.5) hours and 193 (\pm 48) mL/min following oral administration and 16.6 (\pm 5.8) hours and 177 (\pm 31) mL/min following intravenous administration, respectively.

Pharmacokinetics in Healthy Volunteers, Renal, Cardiac and Hepatic Transplant Patients: Shown below are the mean (\pm SD) pharmacokinetic parameters for MPA following the administration of mycophenolate mofetil given as single doses to healthy volunteers and multiple doses to renal, cardiac and hepatic transplant patients. In the early post-transplant period (< 40 days post-transplant), renal, cardiac and hepatic transplant patients had mean MPA AUCs approximately 20% to 41% lower and mean C_{max} approximately 32% to 44% lower compared to the late transplant period (3 to 6 months post-transplant).

Mean MPA AUC values following administration of 1 g bid intravenous mycophenolate mofetil over 2 hours to renal transplant patients for 5 days were about 24% higher than those observed after oral administration of a similar dose in the immediate post-transplant phase. In hepatic transplant patients, administration of 1 g bid intravenous mycophenolate mofetil followed by 1.5 g bid oral mycophenolate mofetil resulted in mean MPA AUC values similar to those found in renal transplant patients administered 1 g mycophenolate mofetil bid.

Table 1 Pharmacokinetic Parameters for MPA (mean \pm SD) Following Administration of Mycophenolate Mofetil to Healthy Volunteers (Single-Dose), Renal, Cardiac and Hepatic Transplant Patients (Multiple Doses)

	Dose/Route	T_{max} (h)	C_{max} (mcg/mL)	Total AUC (mcg•h/mL)
Healthy Volunteers (single-dose)	1 g/oral	0.80 (\pm 0.36) (n = 129)	24.5 (\pm 9.5) (n = 129)	63.9 (\pm 16.2) (n = 117)
Renal Transplant Patients (bid dosing) Time After Transplantation	Dose/Route	T_{max} (h)	C_{max} (mcg/mL)	Interdosing Interval AUC _(0-12h) (mcg•h/mL)
5 days	1 g/iv	1.58 (\pm 0.46) (n = 31)	12 (\pm 3.82) (n = 31)	40.8 (\pm 11.4) (n = 31)
6 days	1 g/oral	1.33 (\pm 1.05) (n = 31)	10.7 (\pm 4.83) (n = 31)	32.9 (\pm 15) (n = 31)
Early (< 40 days)	1 g/oral	1.31 (\pm 0.76) (n = 25)	8.16 (\pm 4.50) (n = 25)	27.3 (\pm 10.9) (n = 25)
Early (< 40 days)	1.5 g/oral	1.21 (\pm 0.81) (n = 27)	13.5 (\pm 8.18) (n = 27)	38.4 (\pm 15.4) (n = 27)
Late (> 3 months)	1.5 g/oral	0.90 (\pm 0.24) (n = 23)	24.1 (\pm 12.1) (n = 23)	65.3 (\pm 35.4) (n = 23)
Cardiac Transplant Patients (bid dosing) Time After Transplantation	Dose/Route	T_{max} (h)	C_{max} (mcg/mL)	Interdosing Interval AUC _(0-12h) (mcg•h/mL)
Early (Day before discharge)	1.5 g/oral	1.8 (\pm 1.3) (n = 11)	11.5 (\pm 6.8) (n = 11)	43.3 (\pm 20.8) (n = 9)
Late (> 6 months)	1.5 g/oral	1.1 (\pm 0.7) (n = 52)	20 (\pm 9.4) (n = 52)	54.1 ^a (\pm 20.4) (n = 49)
Hepatic Transplant Patients (bid dosing) Time After Transplantation	Dose/Route	T_{max} (h)	C_{max} (mcg/mL)	Interdosing Interval AUC _(0-12h) (mcg•h/mL)
4 to 9 days	1 g/iv	1.50 (\pm 0.517) (n = 22)	17 (\pm 12.7) (n = 22)	34 (\pm 17.4) (n = 22)
Early (5 to 8 days)	1.5 g/oral	1.15 (\pm 0.432) (n = 20)	13.1 (\pm 6.76) (n = 20)	29.2 (\pm 11.9) (n = 20)
Late (> 6 months)	1.5 g/oral	1.54 (\pm 0.51) (n = 6)	19.3 (\pm 11.7) (n = 6)	49.3 (\pm 14.8) (n = 6)

^aAUC_(0-12h) values quoted are extrapolated from data from samples collected over 4 hours.

Two 500 mg tablets have been shown to be bioequivalent to four 250 mg capsules. Five mL of the 200 mg/mL constituted oral suspension have been shown to be bioequivalent to four 250 mg capsules.

Special Populations: Shown below are the mean (\pm SD) pharmacokinetic parameters for MPA following the administration of oral mycophenolate mofetil given as single doses to non-transplant subjects with renal or hepatic impairment.

Table 2 Pharmacokinetic Parameters for MPA (mean \pm SD) Following Single Doses of Mycophenolate Mofetil Capsules in Chronic Renal and Hepatic Impairment

Renal Impairment (no. of patients)	Dose	T_{max} (h)	C_{max} (mcg/mL)	AUC _(0-96h) (mcg•h/mL)
Healthy Volunteers	1 g	0.75 (\pm 0.27)	25.3 (\pm 7.99)	45 (\pm 22.6)
Mild Renal Impairment GFR > 80 mL/min/1.73 m ² (n = 6)	1 g	0.75 (\pm 0.27)	26 (\pm 3.82)	59.9 (\pm 12.9)
Moderate Renal Impairment GFR 25 to 49 mL/min/1.73 m ² (n = 6)	1 g	0.75 (\pm 0.27)	19 (\pm 13.2)	52.9 (\pm 25.5)
Severe Renal Impairment GFR < 25 mL/min/1.73 m ² (n = 7)	1 g	1.00 (\pm 0.41)	16.3 (\pm 10.8)	78.6 (\pm 46.4)
Hepatic Impairment (no. of patients)	Dose	T_{max} (h)	C_{max} (mcg/mL)	AUC _(0-48h) (mcg•h/mL)
Healthy Volunteers (n = 6)	1 g	0.63 (\pm 0.14)	24.3 (\pm 5.73)	29 (\pm 5.78)
Alcoholic Cirrhosis (n = 18)	1 g	0.85 (\pm 0.58)	22.4 (\pm 10.1)	29.8 (\pm 10.7)

Renal Insufficiency: In a single-dose study, mycophenolate mofetil was administered as capsule or intravenous infusion over 40 minutes. Plasma MPA AUC observed after oral dosing to volunteers with severe chronic renal impairment [glomerular filtration rate (GFR) < 25 mL/min/1.73 m²] was about 75% higher relative to that observed in healthy volunteers (GFR > 80 mL/min/1.73 m²). In addition, the single-dose plasma MPAG AUC was 3-fold to 6-fold higher in volunteers with severe renal impairment than in volunteers with mild renal impairment or healthy volunteers, consistent with the known renal elimination of MPAG. No data are available on the safety of long-term exposure to this level of MPAG.

Plasma MPA AUC observed after single-dose (1 g) intravenous dosing to volunteers (n = 4) with severe chronic renal impairment (GFR < 25 mL/min/1.73 m²) was 62.4 mcg•h/mL (\pm 19.3). Multiple dosing of mycophenolate mofetil in patients with severe chronic renal impairment has not been studied (see PRECAUTIONS: General and DOSAGE AND ADMINISTRATION).

In patients with delayed renal graft function post-transplant, mean MPA AUC_(0-12h) was comparable to that seen in post-transplant patients without delayed renal graft function. There is a potential for a transient increase in the free fraction and concentration of plasma MPA in patients with delayed renal graft function. However, dose adjustment does not appear to be necessary in patients with delayed renal graft function. Mean plasma MPAG AUC_(0-12h) was 2-fold to 3-fold higher than in post-transplant patients without delayed renal graft function (see PRECAUTIONS: General and DOSAGE AND ADMINISTRATION).

In eight patients with primary graft non-function following renal transplantation, plasma concentrations of MPAG

accumulated about 6-fold to 8-fold after multiple dosing for 28 days. Accumulation of MPA was about 1-fold to 2-fold.

The pharmacokinetics of mycophenolate mofetil are not altered by hemodialysis. Hemodialysis usually does not remove MPA or MPAG. At high concentrations of MPAG (> 100 mcg/mL), hemodialysis removes only small amounts of MPAG.

Hepatic Insufficiency: In a single-dose (1 g oral) study of 18 volunteers with alcoholic cirrhosis and six healthy volunteers, hepatic MPA glucuronidation processes appeared to be relatively unaffected by hepatic parenchymal disease when pharmacokinetic parameters of healthy volunteers and alcoholic cirrhosis patients within this study were compared. However, it should be noted that for unexplained reasons, the healthy volunteers in this study had about a 50% lower AUC as compared to healthy volunteers in other studies, thus making comparisons between volunteers with alcoholic cirrhosis and healthy volunteers difficult. Effects of hepatic disease on this process probably depend on the particular disease. Hepatic disease with other etiologies, such as primary biliary cirrhosis, may show a different effect. In a single-dose (1 g intravenous) study of six volunteers with severe hepatic impairment (aminopyrine breath test less than 0.2% of dose) due to alcoholic cirrhosis, mycophenolate mofetil was rapidly converted to MPA. MPA AUC was 44.1 mcg•h/mL (± 15.5).

Pediatrics: The pharmacokinetic parameters of MPA and MPAG have been evaluated in 55 pediatric patients (ranging from 1 year to 18 years of age) receiving mycophenolate mofetil oral suspension at a dose of 600 mg/m² bid (up to a maximum of 1 g bid) after allogeneic renal transplantation. The pharmacokinetic data for MPA is provided in Table 3:

Table 3 Mean (± SD) Computed Pharmacokinetic Parameters for MPA by Age and Time After Allogeneic Renal Transplantation

Age Group	(n)	Time	T _{max} (h)	Dose Adjusted ^a C _{max} (mcg/mL)	Dose Adjusted ^a AUC ₀₋₁₂ (mcg•h/mL)
Early (Day 7)					
1 to < 2 yr	(6) ^d		3.03 (4.70)	10.3 (5.80)	22.5 (6.66)
1 to < 6 yr	(17)		1.63 (2.85)	13.2 (7.16)	27.4 (9.54)
6 to < 12 yr	(16)		0.940 (0.546)	13.1 (6.30)	33.2 (12.1)
12 to 18 yr	(21)		1.16 (0.830)	11.7 (10.7)	26.3 (9.14) ^b
Late (Month 3)					
1 to < 2 yr	(4) ^d		0.725 (0.276)	23.8 (13.4)	47.4 (14.7)
1 to < 6 yr	(15)		0.989 (0.511)	22.7 (10.1)	49.7 (18.2)
6 to < 12 yr	(14)		1.21 (0.532)	27.8 (14.3)	61.9 (19.6)
12 to 18 yr	(17)		0.978 (0.484)	17.9 (9.57)	53.6 (20.3) ^c
Late (Month 9)					
1 to < 2 yr	(4) ^d		0.604 (0.208)	25.6 (4.25)	55.8 (11.6)
1 to < 6 yr	(12)		0.869 (0.479)	30.4 (9.16)	61 (10.7)
6 to < 12 yr	(11)		1.12 (0.462)	29.2 (12.6)	66.8 (21.2)
12 to 18 yr	(14)		1.09 (0.518)	18.1 (7.29)	56.7 (14)

^a adjusted to a dose of 600 mg/m²

^b n = 20

^c n = 16

^d a subset of 1 to < 6 yr

The mycophenolate mofetil oral suspension dose of 600 mg/m² bid (up to a maximum of 1 g bid) achieved mean MPA AUC values in pediatric patients similar to those seen in adult renal transplant patients receiving mycophenolate mofetil capsules at a dose of 1 g bid in the early post-transplant period. There was wide variability in the data. As observed in adults, early post-transplant MPA AUC values were approximately 45% to 53% lower than those observed in the later post-transplant period (> 3 months). MPA AUC values were similar in the early and late post-transplant period across the 1 year to 18 year age range.

Gender: Data obtained from several studies were pooled to look at any gender-related differences in the pharmacokinetics of MPA (data were adjusted to 1 g oral dose). Mean (± SD) MPA AUC_(0-12h) for males (n = 79) was 32 (± 14.5) and for females (n = 41) was 36.5 (± 18.8) mcg•h/mL while mean (± SD) MPA C_{max} was 9.96 (± 6.19) in the males and 10.6 (± 5.64) mcg/mL in the females. These differences are not of clinical significance.

Geriatrics: Pharmacokinetics in the elderly have not been studied.

CLINICAL STUDIES: Adults: The safety and efficacy of mycophenolate mofetil in combination with corticosteroids and cyclosporine for the prevention of organ rejection were assessed in randomized, double-blind, multicenter trials in renal (three trials), in cardiac (one trial) and in hepatic (one trial) adult transplant patients.

Renal Transplant: Adults: The three renal studies compared two dose levels of oral mycophenolate mofetil (1 g bid and 1.5 g bid) with azathioprine (two studies) or placebo (one study) when administered in combination with cyclosporine (Sandimmune®) and corticosteroids to prevent acute rejection episodes. One study also included antithymocyte globulin (ATGAM®) induction therapy. These studies are described by geographic location of the investigational sites. One study was conducted in the USA at 14 sites, one study was conducted in Europe at 20 sites, and one study was conducted in Europe, Canada and Australia at a total of 21 sites.

The primary efficacy endpoint was the proportion of patients in each treatment group who experienced treatment failure within the first 6 months after transplantation (defined as biopsy-proven acute rejection on treatment or the occurrence of death, graft loss or early termination from the study for any reason without prior biopsy-proven rejection). Mycophenolate mofetil, when administered with antithymocyte globulin (ATGAM®) induction (one study) and with cyclosporine and corticosteroids (all three studies), was compared to the following three therapeutic regimens: (1) antithymocyte globulin (ATGAM®) induction/azathioprine/cyclosporine/corticosteroids, (2) azathioprine/cyclosporine/corticosteroids, and (3) cyclosporine/corticosteroids.

Mycophenolate mofetil, in combination with corticosteroids and cyclosporine reduced (statistically significant at 0.05 level) the incidence of treatment failure within the first 6 months following transplantation. Table 4 and Table 5 summarize the results of these studies. These tables show (1) the proportion of patients experiencing treatment failure, (2) the proportion of patients who experienced biopsy-proven acute rejection on treatment, and (3) early termination, for any reason other than graft loss or death, without a prior biopsy-proven acute rejection episode. Patients who prematurely discontinued treatment were followed for the occurrence of death or graft loss, and the cumulative incidence of graft loss and patient death are summarized separately. Patients who prematurely discontinued treatment were not followed for the occurrence of acute rejection after termination. More patients receiving mycophenolate mofetil discontinued without prior biopsy-proven rejection, death or graft loss than discontinued in the control groups, with the highest rate in the mycophenolate mofetil 3 g/day group. Therefore, the acute rejection rates may be underestimates, particularly in the mycophenolate mofetil 3 g/day group.

Table 4 Renal Transplant Studies Incidence of Treatment Failure (Biopsy-proven Rejection or Early Termination for Any Reason)

USA Study ^a (N = 499 patients)	Mycophenolate Mofetil 2 g/day (n = 167 patients)	Mycophenolate Mofetil 3 g/day (n = 166 patients)	Azathioprine 1 to 2 mg/kg/day (n = 166 patients)
All treatment failures	31.1%	31.3%	47.6%
Early termination without prior acute rejection ^b	9.6%	12.7%	6%
Biopsy-proven rejection episode on treatment	19.8%	17.5%	38%
Europe/Canada/Australia Study ^c (N = 503 patients)	Mycophenolate Mofetil 2 g/day (n = 173 patients)	Mycophenolate Mofetil 3 g/day (n = 164 patients)	Azathioprine 100 to 150 mg/day (n = 166 patients)
All treatment failures	38.2%	34.8%	50%
Early termination without prior acute rejection ^b	13.9%	15.2%	10.2%
Biopsy-proven rejection episode on treatment	19.7%	15.9%	35.5%
Europe Study ^d (N = 491 patients)	Mycophenolate Mofetil 2 g/day (n = 165 patients)	Mycophenolate Mofetil 3 g/day (n = 160 patients)	Placebo (n = 166 patients)
All treatment failures	30.3%	38.8%	56%
Early termination without prior acute rejection ^b	11.5%	22.5%	7.2%
Biopsy-proven rejection episode on treatment	17%	13.8%	46.4%

^a Antithymocyte globulin induction/mycophenolate mofetil or azathioprine/cyclosporine/corticosteroids

^b Does not include death and graft loss as reason for early termination

^c Mycophenolate mofetil or azathioprine/cyclosporine/corticosteroids

^d Mycophenolate mofetil or placebo/cyclosporine/corticosteroids

The cumulative incidence of 12 month graft loss or patient death is presented below. No advantage of mycophenolate mofetil with respect to graft loss or patient death was established. Numerically, patients receiving mycophenolate mofetil 2 g/day and 3 g/day experienced a better outcome than controls in all three studies; patients receiving mycophenolate mofetil 2 g/day experienced a better outcome than mycophenolate mofetil 3 g/day in two of the three studies. Patients in all treatment groups who terminated treatment early were found to have a poor outcome with respect to graft loss or patient death at one year.

Table 5 Renal Transplant Studies Cumulative Incidence of Combined Graft Loss or Patient Death at 12 Months

Study	Mycophenolate Mofetil 2 g/day	Mycophenolate Mofetil 3 g/day	Control (Azathioprine or Placebo)
USA	8.5%	11.5%	12.2%
Europe/Canada/Australia	11.7%	11%	13.6%
Europe	8.5%	10%	11.5%

Pediatrics: One open-label, safety and pharmacokinetic study of mycophenolate mofetil oral suspension 600 mg/m² bid (up to 1 g bid) in combination with cyclosporine and corticosteroids was performed at centers in the US (9), Europe (5) and Australia (1) in 100 pediatric patients (3 months to 18 years of age) for the prevention of renal allograft rejection. Mycophenolate mofetil was well tolerated in pediatric patients (see ADVERSE REACTIONS), and the pharmacokinetics profile was similar to that seen in adult patients dosed with 1 g bid mycophenolate mofetil capsules (see CLINICAL PHARMACOLOGY: Pharmacokinetics). The rate of biopsy-proven rejection was similar across the age groups (3 months to < 6 years, 6 years to < 12 years, 12 years to 18 years). The overall biopsy-proven rejection rate at 6 months was comparable to adults. The combined incidence of graft loss (5%) and patient death (2%) at 12 months post-transplant was similar to that observed in adult renal transplant patients.

Cardiac Transplant: A double-blind, randomized, comparative, parallel-group, multicenter study in primary cardiac transplant recipients was performed at 20 centers in the United States, one in Canada, five in Europe and two in Australia. The total number of patients enrolled was 650; 72 never received study drug and 578 received study drug. Patients received mycophenolate mofetil 1.5 g bid (n = 289) or azathioprine 1.5 to 3 mg/kg/day (n = 289), in combination with cyclosporine (Sandimmune® or Neoral®) and corticosteroids as maintenance immunosuppressive therapy. The two primary efficacy endpoints were: (1) the proportion of patients who, after transplantation, had at least one endomyocardial biopsy-proven rejection with hemodynamic compromise, or were retransplanted or died, within the first 6 months, and (2) the proportion of patients who died or were retransplanted during the first 12 months following transplantation. Patients who prematurely discontinued treatment were followed for the occurrence of allograft rejection for up to 6 months and for the occurrence of death for one year.

(1) **Rejection:** No difference was established between mycophenolate mofetil and azathioprine (AZA) with respect to biopsy-proven rejection with hemodynamic compromise.

(2) **Survival:** Mycophenolate mofetil was shown to be at least as effective as AZA in preventing death or retransplantation at one year (see Table 6).

Table 6 Rejection at 6 Months/Death or Retransplantation at One Year

	All Patients		Treated Patients	
	AZA N = 323	Mycophenolate Mofetil N = 327	AZA N = 289	Mycophenolate Mofetil N = 289
Biopsy-proven rejection with hemodynamic compromise at 6 months ^a	121 (38%)	120 (37%)	100 (35%)	92 (32%)
Death or retransplantation at one year	49 (15.2%)	42 (12.8%)	33 (11.4%)	18 (6.2%)

^a Hemodynamic compromise occurred if any of the following criteria were met: pulmonary capillary wedge pressure ≥ 20 mm or a 25% increase; cardiac index < 2 L/min/m² or a 25% decrease; ejection fraction ≤ 30%; pulmonary artery oxygen saturation ≤ 60% or a 25% decrease; presence of new S₃ gallop; fractional shortening was ≤ 20% or a 25% decrease; inotropic support required to manage the clinical condition.

Hepatic Transplant: A double-blind, randomized, comparative, parallel-group, multicenter study in primary hepatic transplant recipients was performed at 16 centers in the United States, two in Canada, four in Europe and one in Australia. The total number of patients enrolled was 565. Per protocol, patients received mycophenolate mofetil 1 g bid intravenously for up to 14 days followed by mycophenolate mofetil 1.5 g bid orally or azathioprine 1 to 2 mg/kg/day intravenously followed by azathioprine 1 to 2 mg/kg/day orally, in combination with cyclosporine (Neoral®) and corticosteroids as maintenance immunosuppressive therapy. The actual median oral dose of azathioprine on study was 1.5 mg/kg/day (range of 0.3 to 3.8 mg/kg/day) initially and 1.26 mg/kg/day (range of 0.3 to 3.8 mg/kg/day) at 12 months. The two primary endpoints were: (1) the proportion of patients who experienced, in the first 6 months post-transplantation, one or more episodes of biopsy-proven and treated rejection or death or retransplantation, and (2) the proportion of patients who experienced graft loss (death or retransplantation) during the first 12 months post-transplantation. Patients who prematurely discontinued treatment were followed for the occurrence of allograft rejection and for the occurrence of graft loss (death or retransplantation) for one year.

Results: In combination with corticosteroids and cyclosporine, mycophenolate mofetil obtained a lower rate of acute rejection at 6 months and a similar rate of death or retransplantation at one year compared to azathioprine.

Table 7 Rejection at 6 Months/Death or Retransplantation at One Year

	AZA N = 287	Mycophenolate Mofetil N = 278
Biopsy-proven, treated rejection at 6 months (includes death or retransplantation)	137 (47.7%)	107 (38.5%)
Death or retransplantation at one year	42 (14.6%)	41 (14.7%)

INDICATIONS AND USAGE: Renal, Cardiac and Hepatic Transplant: Mycophenolate mofetil is indicated for the prophylaxis of organ rejection in patients receiving allogeneic renal, cardiac or hepatic transplants. Mycophenolate mofetil should be used concomitantly with cyclosporine and corticosteroids.

CONTRAINDICATIONS: Allergic reactions to mycophenolate mofetil have been observed; therefore, mycophenolate mofetil is contraindicated in patients with a hypersensitivity to mycophenolate mofetil, mycophenolic acid or any component of the drug product.

WARNINGS: (see boxed WARNING): Lymphoma and Malignancy: Patients receiving immunosuppressive regimens involving combinations of drugs, including mycophenolate mofetil, as part of an immunosuppressive regimen are at increased risk of developing lymphomas and other malignancies, particularly of the skin (see ADVERSE REACTIONS). The risk appears to be related to the intensity and duration of immunosuppression rather than to the use of any specific agent.

As usual for patients with increased risk for skin cancer, exposure to sunlight and UV light should be limited by wearing protective clothing and using a sunscreen with a high protection factor.

Lymphoproliferative disease or lymphoma developed in 0.4% to 1% of patients receiving mycophenolate mofetil (2 g or 3 g) with other immunosuppressive agents in controlled clinical trials of renal, cardiac and hepatic transplant patients (see ADVERSE REACTIONS).

In pediatric patients, no other malignancies besides lymphoproliferative disorder (2/148 patients) have been observed (see ADVERSE REACTIONS).

Combination with Other Immunosuppressive Agents: Mycophenolate mofetil has been administered in combination with the following agents in clinical trials: antithymocyte globulin (ATGAM®), OKT3 (Orthoclone OKT® 3), cyclosporine (Sandimmune®, Neoral®) and corticosteroids. The efficacy and safety of the use of mycophenolate mofetil in combination with other immunosuppressive agents have not been determined.

Infections: Overtreatment of the immune system can also increase susceptibility to infection, including opportunistic infections, fatal infections and sepsis. In patients receiving mycophenolate mofetil (2 g or 3 g) in controlled studies for prevention of renal, cardiac or hepatic rejection, fatal infection/sepsis occurred in approximately 2% of renal and cardiac patients and in 5% of hepatic patients (see ADVERSE REACTIONS).

Progressive Multifocal Leukoencephalopathy (PML): Cases of progressive multifocal leukoencephalopathy (PML), sometimes fatal, have been reported in patients treated with mycophenolate mofetil. Hemiparesis, apathy, confusion, cognitive deficiencies and ataxia were the most frequent clinical features observed. The reported cases generally had risk factors for PML, including treatment with immunosuppressant therapies and impairment of immune function. In immunosuppressed patients, physicians should consider PML in the differential diagnosis in patients reporting neurological symptoms and consultation with a neurologist should be considered as clinically indicated. Consideration should be given to reducing the amount of immunosuppression in patients who develop PML. In transplant patients, physicians should also consider the risk that reduced immunosuppression represents to the graft.

Pregnancy. Teratogenic Effects. Pregnancy Category D. Mycophenolate mofetil can cause fetal harm when administered to a pregnant woman. Use of mycophenolate mofetil during pregnancy is associated with an increased risk of first trimester pregnancy loss and an increased risk of congenital malformations, especially external ear and other facial abnormalities including cleft lip and palate, and anomalies of the distal limbs, heart, esophagus and kidney. In the National Transplantation Pregnancy Registry (NTPR), there were data on 33 mycophenolate mofetil-exposed pregnancies in 24 transplant patients; there were 15 spontaneous abortions (45%) and 18 live-born infants. Four of these 18 infants had structural malformations (22%). In post-marketing data (collected 1995 to 2007) on 77 women exposed to systemic mycophenolate mofetil during pregnancy, 25 had spontaneous abortions and 14 had a malformed infant or fetus. Six of 14 malformed offspring had ear abnormalities. Because these post-marketing data are reported voluntarily, it is not always possible to reliably estimate the frequency of particular adverse outcomes. These malformations seen in offspring were similar to findings in animal reproductive toxicology studies. For comparison, the background rate for congenital anomalies in the United States is about 3%, and NTPR data show a rate of 4% to 5% among babies born to organ transplant patients using other immunosuppressive drugs.

In animal reproductive toxicology studies, there were increased rates of fetal resorptions and malformations in the absence of maternal toxicity. Female rats and rabbits received mycophenolate mofetil doses equivalent to 0.02 to 0.9 times the recommended human dose for renal and cardiac transplant patients, based on body surface area conversions. In rat offspring, malformations included anophthalmia, agnathia and hydrocephaly. In rabbit offspring, malformations included ectopia cordis, ectopic kidneys, diaphragmatic hernia and umbilical hernia.

If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus. In certain situations, the patient and her healthcare practitioner may decide that the maternal benefits outweigh the risks to the fetus. Women using mycophenolate mofetil at any time during pregnancy should be encouraged to enroll in the National Transplantation Pregnancy Registry.

Pregnancy Exposure Prevention: Women of childbearing potential should have a negative serum or urine pregnancy test with a sensitivity of at least 25 mIU/mL within one week prior to beginning therapy. Mycophenolate mofetil therapy should not be initiated until a negative pregnancy test report is obtained.

Women of childbearing potential (including premenopausal women) taking mycophenolate mofetil must receive contraceptive counseling and use effective contraception. The patient should begin using her two chosen methods of contraception 4 weeks prior to starting mycophenolate mofetil therapy, unless abstinence is the chosen method. She should continue contraceptive use during therapy and for 6 weeks after stopping mycophenolate mofetil. Patients should be aware that mycophenolate mofetil reduces blood levels of the hormones in the oral contraceptive pill and could theoretically reduce its effectiveness (see PRECAUTIONS: Information for Patients and PRECAUTIONS: Drug Interactions: *Oral Contraceptives*).

Neutropenia: Severe neutropenia (absolute neutrophil count (ANC) < 0.5 x 10³/μL) developed in up to 2% of renal, up to 2.8% of cardiac, and up to 3.6% of hepatic transplant patients receiving mycophenolate mofetil 3 g daily (see ADVERSE REACTIONS). Patients receiving mycophenolate mofetil should be monitored for neutropenia (see PRECAUTIONS: Laboratory Tests). The development of neutropenia may be related to mycophenolate mofetil itself, concomitant medications, viral infections, or some combination of these causes. If neutropenia develops (ANC < 1.3 x 10³/μL), dosing with mycophenolate mofetil should be interrupted or the dose reduced, appropriate diagnostic tests performed, and the patient managed appropriately (see DOSAGE AND ADMINISTRATION). Neutropenia has been observed most frequently in the period from 31 to 180 days post-transplant in patients treated for prevention of renal, cardiac and hepatic rejection.

Patients receiving mycophenolate mofetil should be instructed to report immediately any evidence of infection, unexpected bruising, bleeding or any other manifestation of bone marrow depression.

PRECAUTIONS: General: Gastrointestinal bleeding (requiring hospitalization) has been observed in approximately 3% of renal, in 1.7% of cardiac and in 5.4% of hepatic transplant patients treated with mycophenolate mofetil 3 g daily. In pediatric renal transplant patients, 5/148 cases of gastrointestinal bleeding (requiring hospitalization) were observed.

Gastrointestinal perforations have rarely been observed. Most patients receiving mycophenolate mofetil were also receiving other drugs known to be associated with these complications. Patients with active peptic ulcer disease were excluded from enrollment in studies with mycophenolate mofetil. Because mycophenolate mofetil has been associated with an increased incidence of digestive system adverse events, including infrequent cases of gastrointestinal tract ulceration, hemorrhage and perforation, mycophenolate mofetil should be administered with caution in patients with active serious digestive system disease.

Subjects with severe chronic renal impairment (GFR < 25 mL/min/1.73 m²) who have received single doses of mycophenolate mofetil showed higher plasma MPA and MPAG AUCs relative to subjects with lesser degrees of renal impairment or normal healthy volunteers. No data are available on the safety of long-term exposure to these levels of MPAG. Doses of mycophenolate mofetil greater than 1 g administered twice a day to renal transplant patients should be avoided and they should be carefully observed (see CLINICAL PHARMACOLOGY: Pharmacokinetics and DOSAGE AND ADMINISTRATION).

No data are available for cardiac or hepatic transplant patients with severe chronic renal impairment if the potential benefits outweigh the potential risks.

In patients with delayed renal graft function post-transplant, mean MPA AUC_(0-12h) was comparable, but MPAG AUC_(0-12h) was 2-fold to 3-fold higher, compared to that seen in post-transplant patients without delayed renal graft function. In the three controlled studies of prevention of renal rejection, there were 298 of 1,483 patients (20%) with delayed graft function. Although patients with delayed graft function have a higher incidence of certain adverse events (anemia, thrombocytopenia, hyperkalemia) than patients without delayed graft function, these events were not more frequent in patients receiving mycophenolate mofetil than azathioprine or placebo. No dose adjustment is recommended for these patients; however, they should be carefully observed (see CLINICAL PHARMACOLOGY: Pharmacokinetics and DOSAGE AND ADMINISTRATION).

In cardiac transplant patients, the overall incidence of opportunistic infections was approximately 10% higher in patients treated with mycophenolate mofetil than in those receiving azathioprine therapy, but this difference was not associated with excess mortality due to infection/sepsis among patients treated with mycophenolate mofetil (see ADVERSE REACTIONS).

There were more herpes virus (H. simplex, H. zoster and cytomegalovirus) infections in cardiac transplant patients treated with mycophenolate mofetil compared to those treated with azathioprine (see ADVERSE REACTIONS).

It is recommended that mycophenolate mofetil not be administered concomitantly with azathioprine because both have the potential to cause bone marrow suppression and such concomitant administration has not been studied clinically.

In view of the significant reduction in the AUC of MPA by cholestyramine, caution should be used in the concomitant administration of mycophenolate mofetil with drugs that interfere with enterohepatic recirculation because of the potential to reduce the efficacy of mycophenolate mofetil (see PRECAUTIONS: Drug Interactions).

On theoretical grounds, because mycophenolate mofetil is an IMPDH (inosine monophosphate dehydrogenase) inhibitor, it should be avoided in patients with rare hereditary deficiency of hypoxanthine-guanine phosphoribosyl-transferase (HGPRT) such as Lesch-Nyhan and Kelley-Seegmiller syndrome.

During treatment with mycophenolate mofetil, the use of live attenuated vaccines should be avoided and patients should be advised that vaccinations may be less effective (see PRECAUTIONS: Drug Interactions: *Live Vaccines*).

Information for Patients:

- Give patients complete dosage instructions and inform them about the increased risk of lymphoproliferative disease and certain other malignancies.
- Inform patients that they need repeated appropriate laboratory tests while they are taking mycophenolate mofetil.
- Inform women of childbearing potential that use of mycophenolate mofetil in pregnancy is associated with an increased risk of first trimester pregnancy loss and an increased risk of birth defects, and that they must use effective contraception.
- Discuss pregnancy plans with female patients of childbearing potential.
 - Any female of childbearing potential must use highly effective (two methods) contraception 4 weeks prior to starting mycophenolate mofetil therapy and continue contraception until 6 weeks after stopping mycophenolate mofetil treatment, unless abstinence is the chosen method (see WARNINGS: Pregnancy).
 - A patient who is planning a pregnancy should not use mycophenolate mofetil unless she cannot be successfully treated with other immunosuppressant drugs.

Laboratory Tests: Complete blood counts should be performed weekly during the first month, twice monthly for the second and third months of treatment, then monthly through the first year (see WARNINGS, ADVERSE REACTIONS and DOSAGE AND ADMINISTRATION).

Drug Interactions: Drug interaction studies with mycophenolate mofetil have been conducted with acyclovir, antacids, cholestyramine, cyclosporine, ganciclovir, oral contraceptives, sevelamer, trimethoprim/sulfamethoxazole, norfloxacin and metronidazole. Drug interaction studies have not been conducted with other drugs that may be commonly administered to renal, cardiac or hepatic transplant patients. Mycophenolate mofetil has not been administered concomitantly with azathioprine.

Acyclovir: Coadministration of mycophenolate mofetil (1 g) and acyclovir (800 mg) to 12 healthy volunteers resulted in no significant change in MPA AUC and C_{max}. However, MPAG and acyclovir plasma AUCs were increased 10.6% and 21.9%, respectively. Because MPAG plasma concentrations are increased in the presence of renal impairment, as are acyclovir concentrations, the potential exists for mycophenolate and acyclovir or its prodrug (e.g., valacyclovir) to compete for tubular secretion, further increasing the concentrations of both drugs.

Antacids with Magnesium and Aluminum Hydroxides: Absorption of a single-dose of mycophenolate mofetil (2 g) was decreased when administered to ten rheumatoid arthritis patients also taking Maalox[®] TC (10 mL qid). The C_{max} and AUC_(0-24h) for MPA were 33% and 17% lower, respectively, than when mycophenolate mofetil was administered alone under fasting conditions. Mycophenolate mofetil may be administered to patients who are also taking antacids containing magnesium and aluminum hydroxides; however, it is recommended that mycophenolate mofetil and the antacid not be administered simultaneously.

Cholestyramine: Following single-dose administration of 1.5 g mycophenolate mofetil to 12 healthy volunteers pretreated

with 4 g tid of cholestyramine for 4 days, MPA AUC decreased approximately 40%. This decrease is consistent with interruption of enterohepatic recirculation which may be due to binding of recirculating MPAG with cholestyramine in the intestine. Therefore, mycophenolate mofetil is not recommended to be given with cholestyramine or other agents that may interfere with enterohepatic recirculation.

Cyclosporine: Cyclosporine (Sandimmune[®]) pharmacokinetics (at doses of 275 to 415 mg/day) were unaffected by single and multiple doses of 1.5 g bid of mycophenolate mofetil in ten stable renal transplant patients. The mean (± SD) AUC_(0-12h) and C_{max} of cyclosporine after 14 days of multiple doses of mycophenolate mofetil were 3290 (± 822) ng•h/mL and 753 (± 161) ng/mL, respectively, compared to 3245 (± 1088) ng•h/mL and 700 (± 246) ng/mL, respectively, one week before administration of mycophenolate mofetil.

In renal transplant patients, mean MPA exposure (AUC_(0-12h)) was approximately 30% to 50% greater when mycophenolate mofetil is administered without cyclosporine compared with when mycophenolate mofetil is coadministered with cyclosporine. This interaction is due to cyclosporine inhibition of multidrug-resistance-associated protein 2 (MRP-2) transporter in the biliary tract, thereby preventing the excretion of MPAG into the bile that would lead to enterohepatic recirculation of MPA. This information should be taken into consideration when MMF is used without cyclosporine.

Ganciclovir: Following single-dose administration to 12 stable renal transplant patients, no pharmacokinetic interaction was observed between mycophenolate mofetil (1.5 g) and intravenous ganciclovir (5 mg/kg). Mean (± SD) ganciclovir AUC and C_{max} (n = 10) were 54.3 (± 19) mcg•h/mL and 11.5 (± 1.8) mcg/mL, respectively, after coadministration of the two drugs, compared to 51 (± 17) mcg•h/mL and 10.6 (± 2) mcg/mL, respectively, after administration of intravenous ganciclovir alone. The mean (± SD) AUC and C_{max} of MPA (n = 12) after coadministration were 80.9 (± 21.6) mcg•h/mL and 27.8 (± 13.9) mcg/mL, respectively, compared to values of 80.3 (± 16.4) mcg•h/mL and 30.9 (± 11.2) mcg/mL, respectively, after administration of mycophenolate mofetil alone. Because MPAG plasma concentrations are increased in the presence of renal impairment, as are ganciclovir concentrations, the two drugs will compete for tubular secretion and thus further increases in concentrations of both drugs may occur. In patients with renal impairment in which mycophenolate mofetil and ganciclovir or its prodrug (e.g., valganciclovir) are coadministered, patients should be monitored carefully.

Oral Contraceptives: A study of coadministration of mycophenolate mofetil (1 g bid) and combined oral contraceptives containing ethinylestradiol (0.02 mg to 0.04 mg) and levonorgestrel (0.05 mg to 0.20 mg), desogestrel (0.15 mg) or gestodene (0.05 mg to 0.10 mg) was conducted in 18 women with psoriasis over three consecutive menstrual cycles. Mean AUC_(0-24h) was similar for ethinylestradiol and 3-keto desogestrel; however, mean levonorgestrel AUC_(0-24h) significantly decreased by about 15%. There was large inter-patient variability (%CV in the data, especially to 70%) in the data, especially for ethinylestradiol. Mean serum levels of LH, FSH and progesterone were not significantly affected. Mycophenolate mofetil may not have any influence on the ovulation-suppressing action of the studied oral contraceptives. However, it is recommended that oral contraceptives are coadministered with mycophenolate mofetil with caution and additional birth control methods be considered (see WARNINGS: Pregnancy).

Sevelamer: Concomitant administration of sevelamer and mycophenolate mofetil in adult and pediatric patients decreased the mean MPA C_{max} and AUC_(0-12h) by 36% and 26%, respectively. This data suggest that sevelamer and other calcium free phosphate binders should not be administered simultaneously with mycophenolate mofetil. Alternatively, it is recommended that sevelamer and other calcium free phosphate binders preferentially could be given 2 hours after mycophenolate mofetil intake to minimize the impact on the absorption of MPA.

Trimethoprim/Sulfamethoxazole: Following single-dose administration of mycophenolate mofetil (1.5 g) to 12 healthy male volunteers on day 8 of a 10 day course of trimethoprim 160 mg/sulfamethoxazole 800 mg administered bid, no effect on the bioavailability of MPA was observed. The mean (± SD) AUC and C_{max} of MPA after concomitant administration were 75.2 (± 19.8) mcg•h/mL and 34 (± 6.6) mcg/mL, respectively, compared to 79.2 (± 27.9) mcg•h/mL and 34.2 (± 10.7) mcg/mL, respectively, after administration of mycophenolate mofetil alone.

Norfloxacin and Metronidazole: Following single-dose administration of mycophenolate mofetil (1 g) to 11 healthy volunteers on day 4 of a 5 day course of a combination of norfloxacin and metronidazole, the mean MPA AUC_(0-48h) was significantly reduced by 33% compared to the administration of mycophenolate mofetil alone (p < 0.05). Therefore, mycophenolate mofetil is not recommended to be given with the combination of norfloxacin and metronidazole. There was no significant effect on mean MPA AUC_(0-48h) when mycophenolate mofetil was concomitantly administered with norfloxacin or metronidazole separately. The mean (± SD) MPA AUC_(0-48h) after coadministration of mycophenolate mofetil with norfloxacin or metronidazole separately was 48.3 (± 24) mcg•h/mL and 42.7 (± 23) mcg•h/mL, respectively, compared with 56.2 (± 24) mcg•h/mL after administration of mycophenolate mofetil alone.

Rifampin: In a single heart-lung transplant patient, after correction for dose, a 67% decrease in MPA exposure (AUC_(0-12h)) has been observed with concomitant administration of mycophenolate mofetil and rifampin. Therefore, mycophenolate mofetil is not recommended to be given with rifampin concomitantly unless the benefit outweighs the risk.

Other Interactions: The measured value for renal clearance of MPAG indicates removal occurs by renal tubular secretion as well as glomerular filtration. Consistent with this, coadministration of probenecid, a known inhibitor of tubular secretion, with mycophenolate mofetil in monkeys results in a 3-fold increase in plasma MPAG AUC and a 2-fold increase in plasma MPA AUC. Thus, other drugs known to undergo renal tubular secretion may compete with MPAG and thereby raise plasma concentrations of MPAG or the other drug undergoing tubular secretion.

Drugs that alter the gastrointestinal flora may interact with mycophenolate mofetil by disrupting enterohepatic recirculation. Interference of MPAG hydrolysis may lead to less MPA available for absorption.

Live Vaccines: During treatment with mycophenolate mofetil, the use of live attenuated vaccines should be avoided and patients should be advised that vaccinations may be less effective (see PRECAUTIONS: General). Influenza vaccination may be of value. Prescribers should refer to national guidelines for influenza vaccination.

Carcinogenesis, Mutagenesis, Impairment of Fertility: In a 104-week oral carcinogenicity study in mice, mycophenolate mofetil in daily doses up to 180 mg/kg was not tumorigenic. The highest dose tested was 0.5 times the recommended clinical dose (2 g/day) in renal transplant patients and 0.3 times the recommended clinical dose (3 g/day) in cardiac transplant patients when corrected for differences in body surface area (BSA). In a 104-week oral carcinogenicity study in rats, mycophenolate mofetil in daily doses up to 15 mg/kg was not tumorigenic. The highest dose was 0.08 times the recommended clinical dose in renal transplant patients and 0.05 times the recommended clinical dose in cardiac transplant patients when corrected for BSA. While these animal doses were lower than those given to patients, they were maximal in those species and were considered adequate to evaluate the potential for human risk (see WARNINGS).

The genotoxic potential of mycophenolate mofetil was determined in five assays. Mycophenolate mofetil was genotoxic in the mouse lymphoma/thymidine kinase assay and the *in vivo* mouse micronucleus assay. Mycophenolate mofetil was not genotoxic in the bacterial mutation assay, the yeast mitotic gene conversion assay or the Chinese hamster ovary cell chromosomal aberration assay.

Mycophenolate mofetil had no effect on fertility of male rats at oral doses up to 20 mg/kg/day. This dose represents 0.1 times the recommended clinical dose in renal transplant patients and 0.07 times the recommended clinical dose in cardiac transplant patients when corrected for BSA. In a female fertility and reproduction study conducted in rats, oral doses of 4.5 mg/kg/day caused malformations (principally of the head and eyes) in the first generation offspring in the absence of maternal toxicity. This dose was 0.02 times the recommended clinical dose in renal transplant patients and 0.01 times the recommended clinical dose in cardiac transplant patients when corrected for BSA. No effects on fertility or reproductive parameters were evident in the dams or in the subsequent generation.

Pregnancy. Teratogenic Effects. Pregnancy Category D: (See WARNINGS section.)

Nursing Mothers: Studies in rats treated with mycophenolate mofetil have shown mycophenolic acid to be excreted in milk. It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, and because of the potential for serious adverse reactions in nursing infants from mycophenolate mofetil, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use: Based on pharmacokinetic and safety data in pediatric patients after renal transplantation, the recommended dose of mycophenolate mofetil oral suspension is 600 mg/m² bid (up to a maximum of 1 g bid). Also see CLINICAL PHARMACOLOGY, CLINICAL STUDIES, ADVERSE REACTIONS and DOSAGE AND ADMINISTRATION.

Safety and effectiveness in pediatric patients receiving allogeneic cardiac or hepatic transplants have not been established.

Geriatric Use: Clinical studies of mycophenolate mofetil did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal or cardiac function and of concomitant or other drug therapy. Elderly patients may be at an increased risk of adverse reactions compared with younger individuals (see ADVERSE REACTIONS).

ADVERSE REACTIONS: The principal adverse reactions associated with the administration of mycophenolate mofetil include diarrhea, leukopenia, sepsis, vomiting and there is evidence of a higher frequency of certain types of infections e.g., opportunistic infection (see WARNINGS: Infections and WARNINGS: Progressive Multifocal Leukoencephalopathy (PML)).

Mycophenolate Mofetil Oral: The incidence of adverse events for mycophenolate mofetil was determined in randomized, comparative, double-blind trials in prevention of rejection in renal (two active, one placebo-controlled trials), cardiac (one active-controlled trial) and hepatic (one active-controlled trial) transplant patients.

Geriatrics: Elderly patients (> 65 years), particularly those who are receiving mycophenolate mofetil as part of a combination immunosuppressive regimen, may be at increased risk of certain infections (including cytomegalovirus [CMV] tissue invasive disease) and possibly gastrointestinal hemorrhage and pulmonary edema, compared to younger individuals (see PRECAUTIONS).

Safety data are summarized below for all active-controlled trials in renal (two trials), cardiac (one trial) and hepatic (one trial) transplant patients. Approximately 53% of the renal patients, 65% of the cardiac patients and 48% of the hepatic

patients have been treated for more than one year. Adverse events reported in ≥ 20% of patients in the mycophenolate mofetil treatment groups are presented below.

Table 8 Adverse Events in Controlled Studies in Prevention of Renal, Cardiac or Hepatic Allograft Rejection (Reported in ≥ 20% of Patients in the Mycophenolate Mofetil Group)

	Renal Studies			Cardiac Study		Hepatic Study	
	Mycophenolate Mofetil 2 g/day	Mycophenolate Mofetil 3 g/day	Azathioprine 1 to 2 mg/kg/day or 100 to 150 mg/day	Mycophenolate Mofetil 3 g/day	Azathioprine 1.5 to 3 mg/kg/day	Mycophenolate Mofetil 3 g/day	Azathioprine 1 to 2 mg/kg/day
	(n = 336)	(n = 330)	(n = 326)	(n = 289)	(n = 289)	(n = 277)	(n = 287)
	%	%	%	%	%	%	%
Body as a Whole							
Pain	33	31.2	32.2	75.8	74.7	74	77.7
Abdominal pain	24.7	27.6	23	33.9	33.2	62.5	51.2
Fever	21.4	23.3	23.3	47.4	46.4	52.3	56.1
Headache	21.1	16.1	21.2	54.3	51.9	53.8	49.1
Infection	18.2	20.9	19.9	25.6	19.4	27.1	25.1
Sepsis	–	–	–	–	–	27.4	26.5
Asthenia	–	–	–	43.3	36.3	35.4	33.8
Chest pain	–	–	–	26.3	26	–	–
Back pain	–	–	–	34.6	28.4	46.6	47.4
Ascites	–	–	–	–	–	24.2	22.6
Hemic and Lymphatic							
Anemia	25.6	25.8	23.6	42.9	43.9	43	53
Leukopenia	23.2	34.5	24.8	30.4	39.1	45.8	39
Thrombocytopenia	–	–	–	23.5	27	38.3	42.2
Hypochromic anemia	–	–	–	24.6	23.5	–	–
Leukocytosis	–	–	–	40.5	35.6	22.4	21.3
Urogenital							
Urinary tract infection	37.2	37	33.7	–	–	–	–
Kidney function abnormal	–	–	–	21.8	26.3	25.6	28.9
Cardiovascular							
Hypertension	32.4	28.2	32.2	77.5	72.3	62.1	59.6
Hypotension	–	–	–	32.5	36	–	–
Cardiovascular disorder	–	–	–	25.6	24.2	–	–
Tachycardia	–	–	–	20.1	18	22	15.7
Metabolic and Nutritional							
Peripheral edema	28.6	27	28.2	64	53.3	48.4	47.7
Hypercholesterolemia	–	–	–	41.2	38.4	–	–
Edema	–	–	–	26.6	25.6	28.2	28.2
Hypokalemia	–	–	–	31.8	25.6	37.2	41.1
Hyperkalemia	–	–	–	–	–	22	23.7
Hyperglycemia	–	–	–	46.7	52.6	43.7	48.8
Creatinine increased	–	–	–	39.4	36	–	–
BUN increased	–	–	–	34.6	32.5	–	–
Lactic dehydrogenase increased	–	–	–	23.2	17	–	–
Hypomagnesemia	–	–	–	–	–	39	37.6
Hypocalcemia	–	–	–	–	–	30	30
Digestive							
Diarrhea	31	36.1	20.9	45.3	34.3	51.3	49.8
Constipation	22.9	18.5	22.4	41.2	37.7	37.9	38.3
Nausea	19.9	23.6	24.5	54	54.3	54.5	51.2
Dyspepsia	–	–	–	–	–	22.4	20.9
Vomiting	–	–	–	33.9	28.4	32.9	33.4
Anorexia	–	–	–	–	–	25.3	17.1
Liver function tests abnormal	–	–	–	–	–	24.9	19.2
Respiratory							
Infection	22	23.9	19.6	37	35.3	–	–
Dyspnea	–	–	–	36.7	36.3	31	30.3
Cough increased	–	–	–	31.1	25.6	–	–
Lung disorder	–	–	–	30.1	29.1	22	18.8
Sinusitis	–	–	–	26	19	–	–
Pleural effusion	–	–	–	–	–	34.3	35.9
Skin and Appendages							
Rash	–	–	–	22.1	18	–	–
Nervous System							
Tremor	–	–	–	24.2	23.9	33.9	35.5
Insomnia	–	–	–	40.8	37.7	52.3	47
Dizziness	–	–	–	28.7	27.7	–	–
Anxiety	–	–	–	28.4	23.9	–	–
Paresthesia	–	–	–	20.8	18	–	–

The placebo-controlled renal transplant study generally showed fewer adverse events occurring in ≥ 20% of patients. In addition, those that occurred were not only qualitatively similar to the azathioprine-controlled renal transplant studies, but also occurred at lower rates, particularly for infection, leukopenia, hypertension, diarrhea and respiratory infection.

The above data demonstrate that in three controlled trials for prevention of renal rejection, patients receiving 2 g/day of mycophenolate mofetil had an overall better safety profile than did patients receiving 3 g/day of mycophenolate mofetil.

The above data demonstrate that the types of adverse events observed in multicenter controlled trials in renal, cardiac and hepatic transplant patients are qualitatively similar except for those that are unique to the specific organ involved.

Sepsis, which was generally cytomegalovirus viremia, was slightly more common in renal transplant patients treated with mycophenolate mofetil compared to patients treated with azathioprine. The incidence of sepsis was comparable in mycophenolate mofetil and in azathioprine-treated patients in cardiac and hepatic studies.

In the digestive system, diarrhea was increased in renal and cardiac transplant patients receiving mycophenolate mofetil compared to patients receiving azathioprine, but was comparable in hepatic transplant patients treated with mycophenolate mofetil or azathioprine.

Patients receiving mycophenolate mofetil alone or as part of an immunosuppressive regimen are at increased risk of developing lymphomas and other malignancies, particularly of the skin (see WARNINGS: Lymphoma and Malignancy). The incidence of malignancies among the 1,483 patients treated in controlled trials for the prevention of renal allograft rejection who were followed for ≥ one year was similar to the incidence reported in the literature for renal allograft recipients.

Lymphoproliferative disease or lymphoma developed in 0.4% to 1% of patients receiving mycophenolate mofetil (2 g or 3 g daily) with other immunosuppressive agents in controlled clinical trials of renal, cardiac and hepatic transplant patients followed for at least one year (see WARNINGS: Lymphoma and Malignancy). Non-melanoma skin carcinomas occurred in 1.6% to 4.2% of patients, other types of malignancy in 0.7% to 2.1% of patients. Three-year safety data in renal and cardiac transplant patients did not reveal any unexpected changes in incidence of malignancy compared to the one year data.

In pediatric patients, no other malignancies besides lymphoproliferative disorder (2/148 patients) have been observed.

Severe neutropenia (ANC < 0.5 x 10³/μL) developed in up to 2% of renal transplant patients, up to 2.8% of cardiac transplant patients and up to 3.6% of hepatic transplant patients receiving mycophenolate mofetil 3 g daily (see WARNINGS: Neutropenia, PRECAUTIONS: Laboratory Tests and DOSAGE AND ADMINISTRATION).

All transplant patients are at increased risk of opportunistic infections. The risk increases with total immunosuppressive load (see WARNINGS: Infections and WARNINGS: Progressive Multifocal Leukoencephalopathy (PML)). Table 9 shows the incidence of opportunistic infections that occurred in the renal, cardiac and hepatic transplant populations in the azathioprine-controlled prevention trials:

Table 9 Viral and Fungal Infections in Controlled Studies in Prevention of Renal, Cardiac or Hepatic Transplant Rejection

	Renal Studies			Cardiac Study		Hepatic Study	
	Mycophenolate Mofetil 2 g/day	Mycophenolate Mofetil 3 g/day	Azathioprine 1 to 2 mg/kg/day or 100 to 150 mg/day	Mycophenolate Mofetil 3 g/day	Azathioprine 1.5 to 3 mg/kg/day	Mycophenolate Mofetil 3 g/day	Azathioprine 1 to 2 mg/kg/day
	(n = 336)	(n = 330)	(n = 326)	(n = 289)	(n = 289)	(n = 277)	(n = 287)
	%	%	%	%	%	%	%
Herpes simplex	16.7	20	19	20.8	14.5	10.1	5.9
CMV							
– Viremia/syndrome	13.4	12.4	13.8	12.1	10	14.1	12.2
– Tissue invasive disease	8.3	11.5	6.1	11.4	8.7	5.8	8
Herpes zoster	6	7.6	5.8	10.7	5.9	4.3	4.9
– Cutaneous disease	6	7.3	5.5	10	5.5	4.3	4.9
Candida	17	17.3	18.1	18.7	17.6	22.4	24.4
– Mucocutaneous	15.5	16.4	15.3	18	17.3	18.4	17.4

The following other opportunistic infections occurred with an incidence of less than 4% in mycophenolate mofetil patients in the above azathioprine-controlled studies: Herpes zoster, visceral disease; Candida, urinary tract infection, fungemia/dyssemiated disease, tissue invasive disease; Cryptococcosis; Aspergillus/Mucor; Pneumocystis carinii.

In the placebo-controlled renal transplant study, the same pattern of opportunistic infection was observed compared to the azathioprine-controlled renal studies, with a notably lower incidence of the following: Herpes simplex and CMV tissue-invasive disease.

In patients receiving mycophenolate mofetil (2 g or 3 g) in controlled studies for prevention of renal, cardiac or hepatic rejection, fatal infection/sepsis occurred in approximately 2% of renal and cardiac patients and in 5% of hepatic patients (see WARNINGS: Infections).

In cardiac transplant patients, the overall incidence of opportunistic infections was approximately 10% higher in patients treated with mycophenolate mofetil than in those receiving azathioprine, but this difference was not associated with excess mortality due to infection/sepsis among patients treated with mycophenolate mofetil.

The following adverse events were reported with 3% to < 20% incidence in renal, cardiac and hepatic transplant patients treated with mycophenolate mofetil, in combination with cyclosporine and corticosteroids.

Table 10 Adverse Events Reported in 3% to < 20% of Patients Treated With Mycophenolate Mofetil in Combination With Cyclosporine and Corticosteroids

Body System	Adverse Events
Body as a Whole	abdomen enlarged, abscess, accidental injury, cellulitis, chills occurring with fever, cyst, face edema, flu syndrome, hemorrhage, hernia, lab test abnormal, malaise, neck pain, pelvic pain, peritonitis
Hemic and Lymphatic	coagulation disorder, ecchymosis, pancytopenia, petechia, polycythemia, prothrombin time increased, thromboplastin time increased
Urogenital	acute kidney failure, albuminuria, dysuria, hydronephrosis, hematuria, impotence, kidney failure, kidney tubular necrosis, nocturia, oliguria, pain, prostatic disorder, pyelonephritis, scrotal edema, urine abnormality, urinary frequency, urinary incontinence, urinary retention, urinary tract disorder
Cardiovascular	angina pectoris, arrhythmia, arterial thrombosis, atrial fibrillation, atrial flutter, bradycardia, cardiovascular disorder, congestive heart failure, extrasystole, heart arrest, heart failure, hypotension, pallor, palpitation, pericardial effusion, peripheral vascular disorder, postural hypotension, pulmonary hypertension, supraventricular tachycardia, supraventricular extrasystoles, syncope, tachycardia, thrombosis, vasodilation, vasospasm, ventricular extrasystole, ventricular tachycardia, venous pressure increased
Metabolic and Nutritional	abnormal healing, acidosis, alkaline phosphatase increased, alkalosis, bilirubinemia, creatinine increased, dehydration, gamma glutamyl transpeptidase increased, generalized edema, gout, hypercalcemia, hypercholesterolemia, hyperlipemia, hyperphosphatemia, hyperuricemia, hypervolemia, hypocalcemia, hypochloremia, hypoglycemia, hyponatremia, hypophosphatemia, hypoproteinemia, hypovolemia, hypoxia, lactic dehydrogenase increased, respiratory acidosis, SGOT increased, SGPT increased, thirst, weight gain, weight loss
Digestive	anorexia, cholangitis, cholestatic jaundice, dysphagia, esophagitis, flatulence, gastritis, gastroenteritis, gastrointestinal disorder, gastrointestinal hemorrhage, gastrointestinal moniliasis, gingivitis, gum hyperplasia, hepatitis, ileus, infection, jaundice, liver damage, liver function tests abnormal, melena, mouth ulceration, nausea and vomiting, oral moniliasis, rectal disorder, stomach ulcer, stomatitis
Respiratory	apnea, asthma, atelectasis, bronchitis, epistaxis, hemoptysis, hiccup, hyperventilation, lung edema, lung disorder, neoplasm, pain, pharyngitis, pleural effusion, pneumonia, pneumothorax, respiratory disorder, respiratory moniliasis, rhinitis, sinusitis, sputum increased, voice alteration
Skin and Appendages	acne, alopecia, fungal dermatitis, hemorrhage, hirsutism, pruritus, rash, skin benign neoplasm, skin carcinoma, skin disorder, skin hypertrophy, skin ulcer, sweating, vesiculobullous rash
Nervous	agitation, anxiety, confusion, convulsion, delirium, depression, dry mouth, emotional lability, hallucinations, hypertension, hypesthesia, nervousness, neuropraxia, paresthesia, psychosis, somnolence, thinking abnormal, vertigo
Endocrine	Cushing's syndrome, diabetes mellitus, hypothyroidism, parathyroid disorder
Musculoskeletal	arthralgia, joint disorder, leg cramps, myalgia, myasthenia, osteoporosis
Special Senses	abnormal vision, amblyopia, cataract (not specified), conjunctivitis, deafness, ear disorder, ear pain, eye hemorrhage, tinnitus, lacrimation disorder

Pediatrics: The type and frequency of adverse events in a clinical study in 100 pediatric patients 3 months to 18 years of age dosed with mycophenolate mofetil oral suspension 600 mg/m² bid (up to 1 g bid) were generally similar to those observed in adult patients dosed with mycophenolate mofetil capsules at a dose of 1 g bid with the exception of abdominal pain, fever, infection, pain, sepsis, diarrhea, vomiting, pharyngitis, respiratory tract infection, hypertension, leukopenia and anemia, which were observed in a higher proportion in pediatric patients.

Post-Marketing Experience: Congenital Disorders: Congenital malformations including ear malformations have been reported in offspring of patients exposed to mycophenolate mofetil during pregnancy (see WARNINGS: Pregnancy).

Digestive: Colitis (sometimes caused by cytomegalovirus), pancreatitis, isolated cases of intestinal villous atrophy. **Resistance Mechanism Disorders:** Serious life threatening infections such as meningitis and infectious endocarditis have been reported occasionally and there is evidence of a higher frequency of certain types of serious infections such as tuberculosis and atypical mycobacterial infection. Cases of progressive multifocal leukoencephalopathy (PML), sometimes fatal,

have been reported in patients treated with mycophenolate mofetil. The reported cases generally had risk factors for PML, including treatment with immunosuppressant therapies and impairment of immune function.

Respiratory: Interstitial lung disorders, including fatal pulmonary fibrosis, have been reported rarely and should be considered in the differential diagnosis of pulmonary symptoms ranging from dyspnea to respiratory failure in post-transplant patients receiving mycophenolate mofetil.

OVERDOSEAGE: The experience with overdose of mycophenolate mofetil in humans is very limited. The events reported from reports of overdose fall within the known safety profile of the drug. The highest dose administered to renal transplant patients in clinical trials has been 4 g/day. In limited experience with cardiac and hepatic transplant patients in clinical trials, the highest doses used were 4 g/day or 5 g/day. At doses of 4 g/day or 5 g/day, there appears to be a higher rate, compared to the use of 3 g/day or less, of gastrointestinal intolerance (nausea, vomiting and/or diarrhea), and occasional hematologic abnormalities, principally neutropenia, leading to a need to reduce or discontinue dosing.

In acute oral toxicity studies, no deaths occurred in adult mice at doses up to 4000 mg/kg or in adult monkeys at doses up to 1000 mg/kg; these were the highest doses of mycophenolate mofetil tested in these species. These doses represent 11 times the recommended clinical dose in renal transplant patients and approximately 7 times the recommended clinical dose in cardiac transplant patients when corrected for BSA. In adult rats, deaths occurred after single oral doses of 500 mg/kg of mycophenolate mofetil. The dose represents approximately 3 times the recommended clinical dose in cardiac transplant patients when corrected for BSA.

Mycophenolic acid (MPA) and MPAG (metabolized to form the phenolic glucuronide of MPA) are usually not removed by hemodialysis. However, at high MPAG plasma concentrations (> 100 mcg/mL), small amounts of MPAG are removed. By increasing excretion of the drug, MPA can be removed by bile acid sequestrants, such as cholestyramine (see CLINICAL PHARMACOLOGY: Pharmacokinetics).

DOSAGE AND ADMINISTRATION: Renal Transplantation: Adults: A dose of 1 g administered orally twice a day (daily dose of 2 g) is recommended for use in renal transplant patients. Although a dose of 1.5 g administered twice daily (daily dose of 3 g) was used in clinical trials and was shown to be safe and effective, no efficacy advantage could be established for renal transplant patients. Patients receiving 2 g/day of mycophenolate mofetil demonstrated an overall better safety profile than did patients receiving 3 g/day of mycophenolate mofetil.

Pediatrics (3 Months to 18 years of Age): The recommended dose of mycophenolate mofetil oral suspension is 600 mg/m² administered twice daily (up to a maximum daily dose of 2 g/10 mL oral suspension). Patients with a body surface area of 1.25 m² to 1.5 m² may be dosed with mycophenolate mofetil capsules at a dose of 750 mg twice daily (1.5 g daily dose). Patients with a body surface area > 1.5 m² may be dosed with mycophenolate mofetil capsules or tablets at a dose of 1 g twice daily (2 g daily dose).

Cardiac Transplantation: Adults: A dose of 1.5 g bid oral (daily dose of 3 g) is recommended for use in adult cardiac transplant patients.

Hepatic Transplantation: Adults: A dose of 1.5 g bid oral (daily dose of 3 g) is recommended for use in adult hepatic transplant patients.

Mycophenolate Mofetil Capsules and Tablets: The initial oral dose of mycophenolate mofetil should be given as soon as possible following renal, cardiac or hepatic transplantation. Food had no effect on MPA AUC, but has been shown to decrease MPA C_{max} by 40%. Therefore, it is recommended that mycophenolate mofetil be administered on an empty stomach. However, in stable renal transplant patients, mycophenolate mofetil may be administered with food if necessary.

Patients with Hepatic Impairment: No dose adjustments are recommended for renal patients with severe hepatic parenchymal disease. However, it is not known whether dose adjustments are needed for hepatic disease with other etiologies (see CLINICAL PHARMACOLOGY: Pharmacokinetics).

No data are available for cardiac transplant patients with severe hepatic parenchymal disease.

Geriatrics: The recommended oral dose of 1 g bid for renal transplant patients, 1.5 g bid for cardiac transplant patients and 1.5 g bid administered orally in hepatic transplant patients is appropriate for elderly patients (see PRECAUTIONS: Geriatric Use).

Dosage Adjustments: In renal transplant patients with severe chronic renal impairment (GFR < 25 mL/min/1.73 m²) outside the immediate post-transplant period, doses of mycophenolate mofetil greater than 1 g administered twice a day should be avoided. These patients should also be carefully observed. No dose adjustments are needed in renal transplant patients experiencing delayed graft function postoperatively (see CLINICAL PHARMACOLOGY: Pharmacokinetics and PRECAUTIONS: General).

No data are available for cardiac or hepatic transplant patients with severe chronic renal impairment. Mycophenolate mofetil may be used for cardiac or hepatic transplant patients with severe chronic renal impairment if the potential benefits outweigh the potential risks.

If neutropenia develops (ANC < 1.3 x 10³/μL), dosing with mycophenolate mofetil should be interrupted or the dose reduced, appropriate diagnostic tests performed and the patient managed appropriately (see WARNINGS: Neutropenia, ADVERSE REACTIONS and PRECAUTIONS: Laboratory Tests).

HANDLING AND DISPOSAL: Mycophenolate mofetil has demonstrated teratogenic effects in rats and rabbits (see WARNINGS: Pregnancy). Mycophenolate mofetil tablets should not be crushed and mycophenolate mofetil capsules should not be opened or crushed. Avoid inhalation or direct contact with skin or mucous membranes of the powder contained in mycophenolate mofetil capsules. If such contact occurs, wash thoroughly with soap and water; rinse eyes with plain water. Should a spill occur, wipe up using paper towels wetted with water to remove spilled powder.

HOW SUPPLIED: Mycophenolate Mofetil Capsules are available containing 250 mg of mycophenolate mofetil.

The 250 mg capsule is a caramel opaque cap/lavender opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with **MYLAN** over **2250** in black ink on both the cap and body. They are available as follows:

NDC 0378-2250-01
bottles of 100 capsules
NDC 0378-2250-05
bottles of 500 capsules

Mycophenolate Mofetil Tablets are available containing 500 mg of mycophenolate mofetil.

The 500 mg tablet is a light pink film-coated, oval, unscored tablet debossed with **MYLAN** on one side of the tablet and **472** on the other side. They are available as follows:

NDC 0378-4472-01
bottles of 100 tablets
NDC 0378-4472-05
bottles of 500 tablets

Store at 20° to 25°C (68° to 77°F). [See USP for Controlled Room Temperature.]

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

PHARMACIST: Dispense a Medication Guide with each prescription.

Sandimmune® is a registered trademark of Novartis Pharmaceuticals Corporation.

ATGAM® is a registered trademark of Pharmacia and Upjohn Company.

Neoral® is a registered trademark of Novartis Pharmaceuticals Corporation.

Orthoclone OKT® is a registered trademark of Ortho Biotech Inc.

Maalox® is a registered trademark of Novartis Consumer Health, Inc.



Mylan Pharmaceuticals Inc.
Morgantown, WV 26505

REVISED FEBRUARY 2009
MYCO-R2mpbmt

MEDICATION GUIDE

MYCOPHENOLATE MOFETIL CAPSULES and MYCOPHENOLATE MOFETIL TABLETS

Read the Medication Guide that comes with mycophenolate mofetil capsules and mycophenolate mofetil tablets before you start taking it and each time you refill your prescription. There may be new information. This Medication Guide does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about mycophenolate mofetil?

Mycophenolate mofetil can cause serious side effects:

- **Possible loss of a pregnancy and higher risk of birth defects.** Women who take mycophenolate mofetil during pregnancy have a higher risk of losing a pregnancy (miscarriage) during the first 3 months (first trimester), and a higher risk that their baby will be born with birth defects.
 - If you are a female and are able to become pregnant:
 - Your healthcare provider must talk with you about effective birth control methods (contraceptive counseling).
 - You should have a negative pregnancy test within one week before you start to take mycophenolate mofetil.
 - You must use two different types of effective birth control at the same time, for 4 weeks before you start taking mycophenolate mofetil, during your entire mycophenolate mofetil therapy and for 6 weeks after stopping mycophenolate mofetil, unless you choose to avoid sexual intercourse completely (abstinence). Mycophenolate mofetil decreases blood levels of the hormones in birth control pills that you take by mouth. Birth control pills may not work as well while you take mycophenolate mofetil, and you could become pregnant.
 - If you plan to become pregnant, talk with your healthcare provider. Your healthcare provider will decide if other medicines to prevent rejection may be right for you. In certain situations, you and your healthcare provider may decide that taking mycophenolate mofetil is more important to your health than the possible risks to your unborn baby.
- **If you get pregnant while taking mycophenolate mofetil, do not stop taking mycophenolate mofetil. Call your healthcare provider right away.** You and your healthcare provider should report any cases of pregnancies to:
 - FDA MedWatch at 1-800-FDA-1088
 - Mylan Pharmaceuticals Inc. at 1-877-446-3679 (1-877-4-INFO-RX)

Talk to your healthcare provider about joining the National Transplantation Pregnancy Registry at 1-877-955-6877.

- **Increased risk of getting serious infections.** Mycophenolate mofetil weakens the body's immune system and affects your ability to fight infections. Serious infections can happen with mycophenolate mofetil and can lead to death. Types of infections can include:
 - **Viral infections.** Viral infections, including shingles, other herpes infections, and cytomegalovirus (CMV), can happen with mycophenolate mofetil. CMV can cause serious tissue and blood infections.
 - **A brain infection called Progressive Multifocal Leukoencephalopathy (PML).** In some patients, mycophenolate mofetil may cause an infection of the brain that may cause death. You are at risk for this brain infection because you have a weakened immune system. You should tell your healthcare provider right away if you have any of the following symptoms:
 - Weakness on one side of the body
 - You do not care about things that you usually care about (apathy).
 - You are confused or have problems thinking.
 - You can not control your muscles.
 - **Fungal infections.** Yeasts and other types of fungal infections can happen with mycophenolate mofetil and can cause serious tissue and blood infections (see "What are the possible side effects of mycophenolate mofetil?").

Call your healthcare provider right away if you have any of the following signs and symptoms of infection:

- Temperature of 100.5°F or greater
- Cold symptoms, such as a runny nose or sore throat
- Flu symptoms, such as an upset stomach, stomach pain, vomiting or diarrhea
- Earache or headache
- Pain during urination
- White patches in the mouth or throat
- Unexpected bruising or bleeding
- Cuts, scrapes or incisions that are red, warm and oozing pus
- **Increased risk of getting certain cancers.** People who take mycophenolate mofetil have a higher risk of getting lymphoma, and other cancers, especially skin cancer. Tell your healthcare provider if you have:
 - unexplained fever, prolonged tiredness, weight loss or lymph node swelling
 - a brown or black skin lesion with uneven borders, or one part of the lesion does not look like the other
 - a change in the size and color of a mole
 - a new skin lesion or bump
 - any other changes to your health

See the section "What are the possible side effects of mycophenolate mofetil?" for information about other serious side effects.

What is mycophenolate mofetil?

Mycophenolate mofetil is a prescription medicine to prevent rejection (antirejection medicine) in people who have received a kidney, heart or liver transplant. Rejection is when the body's immune system perceives the new organ as a "foreign" threat and attacks it.

Mycophenolate mofetil is used with other medicines called cyclosporines (Sandimmune®, Gengraf®, Neoral®) and corticosteroids. These medicines work together to prevent rejection to your transplanted organ.

Mycophenolate mofetil has been used safely and works in children who received a kidney transplant as it does in adults. It is not known if mycophenolate mofetil is safe and works in children who receive a heart or liver transplant.

Who should not take mycophenolate mofetil?

Do not take mycophenolate mofetil if you are allergic to mycophenolate mofetil or any of the ingredients in mycophenolate mofetil capsules and tablets. See the end of this Medication Guide for a complete list of ingredients in mycophenolate mofetil capsules and tablets.

What should I tell my healthcare provider before taking mycophenolate mofetil?

Tell your healthcare provider about all of your medical conditions, if you:

- have any digestive problems, such as ulcers
- have Lesch-Nyhan or Kelley-Seegmiller syndrome or another rare inherited deficiency hypoxanthine-guanine phosphoribosyl-transferase (HGPRT). You should not take mycophenolate mofetil if you have one of these disorders.
- plan to receive any vaccines. **People taking mycophenolate mofetil should not take live vaccines.** Some vaccines may not work as well during treatment with mycophenolate mofetil.
- are pregnant or are planning to become pregnant. See "What is the most important information I should know about mycophenolate mofetil?"
- are breast-feeding. It is not known if mycophenolate mofetil passes into breast milk. You and your healthcare provider will decide if you will take mycophenolate mofetil or breast-feed. You should not do both without first talking with your healthcare provider.

Tell your healthcare provider about all of the medicines you are taking including prescription and nonprescription medicines, vitamins and herbal supplements. Some medicines may affect the way mycophenolate mofetil works, and mycophenolate mofetil may affect how some medicines work. Especially tell your healthcare provider if you take:

- birth control pills (oral contraceptives). See "What is the most important information I should know about mycophenolate mofetil?"
- sevelamer (Renagel®), Renvela™). These products should be taken 2 hours after taking mycophenolate mofetil.
- acyclovir (Zovirax®), valacyclovir (Valtrex®), ganciclovir (Cytovene®-IV, Valcyte®)
- rifampin (Rifater®, Rifamate®, Rimactan®, Rifadin®)
- antacids that contain magnesium and aluminum (mycophenolate mofetil and the antacid should not be taken at the same time)
- sulfamethoxazole/trimethoprim (Bactrim™, Bactrim DST™)
- norfloxacin (Noroxin®) and metronidazole (Flagyl® ER, Flagyl® IV, Metro® IV, Helidac®, Pylera™)
- azathioprine (Azasan®, Imuran®)
- cholestyramine (Questran Light®, Questran®, Locholest Light®, Locholest®, Prevalite®)

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine. Do not take any new medicine without talking with your healthcare provider.

How should I take mycophenolate mofetil?

- Take mycophenolate mofetil exactly as prescribed.
- Do not stop taking mycophenolate mofetil or change the dose unless your healthcare provider tells you to.
- If you miss a dose of mycophenolate mofetil, or are not sure when you took your last dose, take the regular amount of mycophenolate mofetil prescribed as soon as you remember. If it is time for your next dose, skip the missed dose. Do not take two doses at the same time. Call your healthcare provider if you are not sure what to do.
- Take mycophenolate mofetil capsules and tablets on an empty stomach, either 1 hour before or 2 hours after a meal, unless your healthcare provider tells you otherwise. With the approval of your healthcare provider, in stable kidney transplant patients, mycophenolate mofetil can be taken with food if necessary.
- Most people take mycophenolate mofetil by mouth either as caramel and lavender capsules or light pink tablets. Some people may get mycophenolate mofetil soon after their transplant surgery as an infusion into a vein.
- Do not crush mycophenolate mofetil tablets. Do not open or crush mycophenolate mofetil capsules.
- If you are not able to swallow mycophenolate mofetil tablets or capsules, your healthcare provider may prescribe mycophenolate mofetil oral suspension. This is a liquid form of mycophenolate mofetil. Your pharmacist will mix the medicine before giving it to you.
- Do not mix mycophenolate mofetil oral suspension with any other medicine.
- If you take too much mycophenolate mofetil, call your healthcare provider or the poison control center right away.

What should I avoid while taking mycophenolate mofetil?

- **Avoid pregnancy.** See "**What is the most important information I should know about mycophenolate mofetil?**".
- **Limit the amount of time you spend in sunlight.** Avoid using tanning beds or sunlamps. People who take mycophenolate mofetil have a higher risk of getting skin cancer. (See "**What is the most important information I should know about mycophenolate mofetil?**".) Wear protective clothing when you are in the sun and use a sunscreen with a high protection factor (SPF 30 and above). This is especially important if your skin is very fair or if you have a family history of skin cancer.

What are the possible side effects of mycophenolate mofetil?

Mycophenolate mofetil can cause serious side effects:

- See "**What is the most important information I should know about mycophenolate mofetil?**".
- **Low blood cell counts.** People taking high doses of mycophenolate mofetil each day may have a decrease in blood counts, including:
 - **white blood cells, especially neutrophils.** Neutrophils fight against bacterial infections. You have a higher chance of getting an infection when your white blood cell count is low. This is most common from 3 months to 6 months after your transplant.
 - **red blood cells.** Red blood cells carry oxygen to your body tissues.
 - **platelets.** Platelets help with blood clotting.Your healthcare provider will do blood tests before you start taking mycophenolate mofetil and during treatment with mycophenolate mofetil to check your blood cell counts. Tell your healthcare provider right away if you have any signs of infection (see "**What is the most important information I should know about mycophenolate mofetil?**"). Also, tell your healthcare provider about any unexpected bruising or bleeding.
- **Stomach problems.** Stomach and intestinal bleeding can happen in people who take high doses of mycophenolate mofetil. Bleeding can be severe and you may have to be hospitalized for treatment.

Common side effects include:

- diarrhea. Call your healthcare provider right away if you have diarrhea. Do not stop taking mycophenolate mofetil without first talking with your healthcare provider.
- vomiting
- pain
- stomach area pain
- swelling of the lower legs, ankles and feet
- high blood pressure

Side effects that happen more often in children than in adults taking mycophenolate mofetil include:

- stomach area pain
- fever
- infection
- pain
- blood infection (sepsis)
- diarrhea
- vomiting
- sore throat
- colds (respiratory tract infections)
- high blood pressure
- low white blood cell count
- low red blood cell count

These are not all of the possible side effects of mycophenolate mofetil. Tell your healthcare provider about any side effect that bothers you or that does not go away.

How should I store mycophenolate mofetil?

- Store mycophenolate mofetil capsules and tablets at room temperature 20° to 25°C (68° to 77°F). Keep the container closed tightly.
- **Keep mycophenolate mofetil and all medicines out of the reach of children.**

General information about mycophenolate mofetil:

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use mycophenolate mofetil for a condition for which it was not prescribed. Do not give mycophenolate mofetil to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about mycophenolate mofetil. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about mycophenolate mofetil that is written for healthcare professionals. For more information call Mylan Pharmaceuticals Inc. at 1-877-446-3679 (1-877-4-INFO-RX).

What are the ingredients in mycophenolate mofetil capsules?

Active Ingredient: mycophenolate mofetil

Inactive Ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, pregelatinized starch, and sodium lauryl sulfate. The empty gelatin capsule shells contain black iron oxide, FD&C Blue No. 2, gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. In addition, the imprinting ink contains the following: ammonium hydroxide, black iron oxide, propylene glycol, and shellac glaze.

What are the ingredients in mycophenolate mofetil tablets?

Active Ingredient: mycophenolate mofetil

Inactive Ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, red iron oxide, sodium lauryl sulfate, talc, titanium dioxide and yellow iron oxide.

† The brand names mentioned in this Medication Guide are registered trademarks of their respective owners.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

This Medication Guide has been approved by the US Food and Drug Administration.



MYLAN®

Mylan Pharmaceuticals Inc.
Morgantown, WV 26505

FEBRUARY 2009
MG:MYCO:R1